

Shri X-Ray & Ultrasound Centre

Equipped with DIGITAL X-RAYS & 300 mA Xray Plant & Latest JAPANESE
TOSIBA WHOLE BODY COLOR DOPPLER ULTRASOUND SCAN
Main Jagadhri Road, Mahesh Nagar, Ambala Cantt. - 133 001

DR. MUKESH GUPTA

M.B.B.S., M.D. (RADIODIAGNOSIS)

Consultant Radiologist & Ultrasonologist

Timings : Morning : 9-00 A.M. to 1-30 P.M.

Evening : 5-00 P.M. to 7-30 P.M.

DR. SUMIT GUPTA

M.B.B.S., M.D. (RADIODIAGNOSIS)

Ex. Consultant Radiologist & Ultrasonologist

Rotary Ambala Cancer & General Hospital

Timing : 5-00 P.M. to 8-00 P.M.

PATIENT NAME: [REDACTED]

M

13-04-2023

X- RAY CHEST PA :

REPORT :

- 1] Heart size is Normal .
- 2] Lung Fields are clear .
- 3] Both costophrenic Angles are clear .
- 4] Thoracic cage is Normal .

CONCLUSION : Xray chest PA : Normal study .

^{MG}
DR MUKESH GUPTA
M.D. [RADIODIAGNOSIS]

Col & M.D.

P.D. Sharma

Col (Dr.) P.D. SHARMA
M.B.B.S., M.D. (PAED)
CONSULTANT PEDIATRICIAN
REGN No. MCI : 18-27302



REG NO. : 2,731	
PATIENT NAME : [REDACTED]	UHID NUMBER : [REDACTED]
AGE/ GENDER : 18 Year/Male	REGISTRATION DATE : 13-04-2023 01:53:00 PM
REFERRED BY DR.: SELF	COLLECTION DATE : 13-04-2023 01:55 PM
SPECIMEN TYPE : BLOOD	REPORTING DATE : 13-04-2023 02:22:07 PM

Investigations	Result	Unit	Biological Reference Range
----------------	--------	------	----------------------------

SEROLOGY

HEPATITIS C VIRUS NON REACTIVE NON REACTIVE

Comments:

HEPATITIS C VIRUS accounts for about 95% of hepatitis infections in recipients of blood transfusion and 50% of cases of Sporadic nanb hepatitis. HCV commonly gives origin to asymptomatic hepatitis and chronicity develop in a high number of cases, sometime evolving in severs forms of illness, as hepatocarcinoma. IGM Antibodies directed to the major immunodominant patient determinants of the viral proteins are detected in patients infected with HCV, early in the courses of infection and in patients upon reactivation of viral relocation in Hepatocytes.

HEPATITIS B SURFACE ANTIGEN NON REACTIVE NON REACTIVE

COMMENT : This assay detects the first serological marker of Hepatitis B as early as 4-16 weeks after exposure. It persists during acute illness and disappears 12-20 week after onset of symptoms. The titres during the period of viral replication and is frequently associated with infectivity. Persistence of HBsAg for more than 6 months indicates development of carrier state or chronic liver disease. Uses Routine screening of blood and blood products to prevent prevent transmission of Hepatitis B virus (HBS) to recipients. To diagnose suspected HBV infection and monitor the status of infected individuals To evaluate the efficacy of antiviral drugs. For prenatal Screening of pregnant women

V.D.R.L. NON REACTIVE NON REACTIVE

COMMENTS: VDRL detects antibodies found is early syphilis, but can be nonreactive is later stages Biologic false positives are common in a variety of other infections or rheumatic diseases. Repeat samples after two weeks. More specific T.pallidum tests are recommended for confirmation

HIV 1 & 2 NON REACTIVE NON REACTIVE

Dr. RAVINDER PAL
Ph.D. (Bio Chemistry PGIMER CHD)
Pst DOC (USA & Canada)
Regd. No. 6231-PG-2001

Col. (Dr.) P.D. SHARMA
M.D. (PAEDI)
CONSULTANT PEDIATRICIAN
REGN NO. MCI : 18-27302

A. K. SAINI
M.Sc. - MLT
LAB TECHNOLOGIST



GANESH LAB

FULLY COMPUTERISED LABORATORY

43 - Palika Bazar, Opp. Local Bus Stand, Mahesh Nagar, Ambala Cantt.
Mb. 94162 24201, 92152 24201 Ph. 0171-4000220



e-reports

Save paper- Save tree- Save the environment

Email : shriganeshlab201@gmail.com

REG NO. : 2,731



PATIENT NAME : [REDACTED]

UHID NUMBER [REDACTED]

AGE/ GENDER : 18 Year/Male

REGISTRATION DATE : 13-04-2023 01:53:00 PM

REFERRED BY DR.: SELF

COLLECTION DATE : 13-04-2023 01:55 PM


SPECIMEN TYPE : BLOOD

REPORTING DATE : 13-04-2023 02:22:07 PM

Investigations	Result	Unit	Biological Reference Range
----------------	--------	------	----------------------------

COMMENTS: Non reactive results indicate that antibodies to HIV I/II have not been detected in the sample. This implies either non exposure to HIV I/II or that the individual might be in the 'Window Period', that is, prior to the development of detectable level of Antibodies. Therefore, a non-reactive result does not exclude the possibility of a persisting HIV infection. Reactive samples must be confirmed by using HIV Western Blot as some degree of cross reactivity for HIV antibodies has been noted with certain other naturally occurring antibodies and also antibodies formed in response to other bacterial and viral infections, and in certain autoimmune disorders. Post test counseling will be done by the concerned referring doctor. The sensitivity and specificity of this test has been determined by National HIV Reference Centers of Govt .of India and WHO collaborating cent

-----End of Report-----



[Handwritten signature]

Dr. RAVINDER PAL
Ph.D. (Bio Chemistry PGIMER CHD)
Pst DOC (USA & Canada)
Regd. No. 6231-PG-2001

[Handwritten signature]

Col (Dr.) P.D. SHARMA
M.D. (PAED) 11/1/23
CONSULTANT PEDIATRICIAN
REGN NO. MCI : 18-27302

[Handwritten signature]

A. K. SAINI
M.Sc. - MLT
LAB TECHNOLOGIST



Authorized Collection Centre : FAMILY HEALTH CARE CENTRE
43, 1st Floor, Palika Bazar, Near Sethi Banquet Hall, Mahesh Nagar, Ambala Cantt.